



Divergence Study of Rice (*Oryza sativa* L) Genotypes for Consumer-preferred Quality Traits

Amrutlal R. Khaire ^{a*}, S. K. Singh ^a, M. Jayasudha ^a, Akanksha Singh ^a,
D. K. Singh ^{a,b}, Korada Mounika ^a, Sonali Vijay Habde ^a,
Prasanta Kumar Manzhi ^a, Arsode Pandurang Bhagvan ^a
and Saurav Singla ^c

^a Department of Genetics and Plant Breeding, Institute of Agricultural Sciences, Banaras Hindu University, Varanasi, Uttar Pradesh -221005, India.

^b Department of Genetics and Plant Breeding, College of Horticulture, Banda University of Agriculture and Technology, Banda - 210001, Uttar Pradesh, India.

^c Department of Farm Engineering, Institute of Agricultural Sciences, Banaras Hindu University, Varanasi, Uttar Pradesh -221005, India.

Authors' contributions

This work was carried out in collaboration among all authors. Authors ARK, SKS and MJ designed the study and wrote the protocol. Authors, ARK, AS, SS and KM performed the statistical analysis and wrote the first draft of manuscript. Authors DKS, SVH, APB and PKM managed the further analyses and improvement in the manuscript and also, managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JEAI/2022/v44i112061

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/92540>

Original Research Article

Received 05 August 2022

Accepted 09 October 2022

Published 13 October 2022

ABSTRACT

Aim: To study the divergence of forty rice genotypes for consumer-preferred eleven quality traits.
Design, Place and Duration of the Study: The rice seeds were harvested from randomly selected plants grown in alpha lattice design with three replications during *Kharif* season (start in June and end in October) 2018 at Agricultural research farm, Banaras Hindu University, Varanasi UP, India.
Methodology: The data were analysed using biometrical tools - Mahalanobis D² and Principal Component Analysis (PCA). The studied traits include, - grain length, grain breadth, kernel length, length to breadth ratio, kernel breadth, kernel length after cooking, kernel breadth after cooking, elongation ratio, elongation index, amylose content and alkali digestion value.

*Corresponding author: E-mail: amrutlalrkhair10@bhu.ac.in;

Results: Using Mahalanobis D^2 , all forty genotypes were distributed into six clusters. The maximum inter-cluster distance was recorded between Cluster II to cluster V (5.76), followed by cluster V and cluster III (5.71), and cluster II and cluster VI (5.57) which indicated the existence of high genetic diversity among genotypes in these clusters and, therefore, crosses between the genotypes of these clusters could yield desirable transgressive isolates for desirable quality traits and the importance of the genotypes present in these clusters for exploiting heterosis for the desirable traits of these clusters. The PCA indicates that the five principal components (PC) captured almost 90% of variability present among the 40 rice genotypes.

Conclusion: The genotypes belonging to Cluster II to cluster V, followed by cluster V and cluster III, and cluster II and cluster VI, can be used for making crosses as they have higher mean values for quality traits and higher inter-cluster distance for greater diversity.

Keywords: Genetic divergence; rice; quality traits; principal component; Mahalanobis D^2 .

1. INTRODUCTION

"Rice (*Oryza sativa* L.) is the most important crop in the world as it is present in the daily dishes of two-thirds of the world's population" [1]. Developing the variety with better yield is one of the foremost vital goals of rice breeders. The adaptability of any variety becomes crucial if it has significant rice quality traits. The consumer market has recently seen an increase in quality sensitivity. According to a survey of 11 major rice-growing nations, grain quality is the second-most important breeding goal behind production [2]. Various rice products are always in demand by both, poor to rich people, and the requirement of quality accords with country, continent, state, and even person to person.

Grain dimension quality traits i.e. kernel length, kernel breadth, length to breadth ratio, elongation ratio, kernel length after cooking, and kernel breadth after cooking have great importance in commercial rice production as it highly influences one final output as well as the consumer demand which directly contributes the economic profitability of the rice cultivator.

The Physicochemical properties of rice are determined based on amylose content, gel consistency and gelatinisation which depends on cooking and eating qualities and greatly influences the consumer's affinity [3,4]. Amylose content determines the texture of cooked rice. The term "resistant starch" refers to amylose, composed of linearly connected glucose molecules and relatively difficult to digest. As a result, rice types with higher amylose starch content tend to have lower glycemic indexes. It has been discovered that there is positive correlation between the amylose content of milled rice and hardness values of cooked rice, and negative correlation between the amylose

content and stickiness values. The intermediate range of amylose content (21-25%) is desirable in the Indian population. The eating and cooking quality of rice grains can therefore be considered a significant critical quality component that must be prioritized in the country's future rice breeding programs to meet market needs worldwide.

"To achieve this objective, the breeder must identify diverse parents with high genetic variability for combining desirable characters. Therefore, the knowledge of genetic diversity is essential for undertaking any recombination breeding program. Multivariate statistical techniques that assess several variables on each individual under investigation are frequently used to genetic diversity. Principal component analysis (PCA) and cluster analysis have been demonstrated to be particularly helpful in selecting genotypes for breeding programs that satisfy a plant breeder's objectives" [5].

This study was carried out to evaluate the variety of rice grain quality, various genotypes on physical and physicochemical qualities. The results of this study will be hugely beneficial for both, customers and future rice breeding operations.

2. MATERIALS AND METHODS

The material of the present experiment consists of forty rice genotypes sown at Agricultural Research Farm, Banaras Hindu University, Varanasi, Uttar Pradesh, India during Kharif - 2018. The 25 days old seedlings were transplanted in Alpha Lattice design by adopting the spacing of 25 x 15 cm and recommended package of practices, and following all plant protection measures. At harvest maturity stage, with the moisture content of 12-14%, the observed seed quality traits were recorded. The

samples from each replication were taken into consideration.

2.1 Observation Recorded

The following physical and physiochemical observations were recorded:

1. **Grain length (GL):** The length of 10 grains from its base to tip was recorded (mm).
2. **Grain breadth (GB):** Grain breadth was measured at the widest point of the 10 grains (mm)
3. **Kernel length (KL):** Length of 10 dehulled (after milling) grains from its base to tip was recorded (mm).
4. **Kernel breadth (KB):** Kernel breadth was measured at the widest point of the 10 kernels (mm).
5. **L/B ratio (LBBC):** Value was obtained by dividing kernel length by kernel breadth before cooking.
6. **Kernel length after cooking (KLAC):** Kernel length after cooking for 10 minutes in a water bath was recorded (mm).
7. **Kernel breadth after cooking (KBAC):** Kernel breadth after cooking for 10 minutes in a water bath was recorded (mm).
8. **Elongation ratio (ER):** Kernel length (mm) of 10 grains after cooking was divided by kernel length (mm) before cooking.
9. **Elongation Index (EI):** It was obtained by dividing by L/B ratio of the cooked kernel by L/B ratio of the raw kernel.
10. **Alkali digestion value / Gelatinization Temperature (ADV/GT):** The spread of the 6 milled rice kernels in 1.7 percent KOH solution for 23 hours was rated as per the Standard Evaluation System for Rice (IRRI, 1996). Six milled kernels were placed in 10 ml of 1.7 percent KOH solution in a Petri dish and arranged in a manner not to touch each other and were allowed to stand for 23 hours at 30 °C to score spreading on a 1-7 scale.
11. **Amylose content (Amylcnt):** Amylose content (%) was calculated using the simplified procedure described by Juliano [6]. In a 100 ml volumetric flask, 100 mg of ground milled rice sample (sieved with a mesh size of 60) was placed. One ml of 95% ethanol and 9 ml of 1N sodium hydroxide were added. After being heated in a boiling water bath to gelatinize the starch, the mixture was cooled for an hour before being thoroughly mixed with distilled

water up to the volume. Starch solution (5 ml), 1 ml of 1N acetic acid, and 2 ml of iodine solution were added to a 100 ml volumetric flask and the volume was made with distilled water. After thoroughly shaking the contents, let them stand for 20 minutes and then absorbance was taken at 620 nm using a spectrometer.

2.2 Statistical Analysis

Mahalanobis D^2 values using Tocher's method was used to carry out analysis of forty rice genotypes for all 11 characters, using the mean values of all recorded attributes from all genotypes in all replications Rao [7]. According to Singh and Choudhary [8], the D^2 values for each combination were arranged in a table in ascending order. Each character was ranked according to how much they contributed to the divergence between the two entries. For every combination of entries, each character was ranked from rank "1" (character with a highest mean difference) to rank "p." (character with the lowest mean difference).

3. RESULT AND DISCUSSION

3.1 Cluster Analysis

The study of genetic diversity and the creation of core subsets for classifying accessions with comparable characteristics into a single homogenous category are two common uses for cluster analysis. By combining related units, clustering can also be utilized to communicate and summarize information about the relationships between various things. In the present experiment, "forty genotypes were grouped into six clusters using Tocher's method" [9] based on Mahalanobis D^2 values. Clusters with their genotypes are presented in table 1. Cluster I had nine genotypes, Cluster II had five genotypes, Cluster III had thirteen, cluster IV had seven genotypes, cluster V had two genotypes, VI had 4 genotypes.

Range of average intra-cluster D^2 values was 0.77 to 1.32 (Table 2). Maximum intra cluster distance was shown by cluster V (1.32). While, cluster IV had minimum intra cluster value (0.77). Intra cluster values of 1.26, 1.07, 1.06, 0.92 were possessed by cluster I, cluster II, cluster III, cluster V and cluster VI respectively. The cluster V had high intra-cluster distance indicate wide diversity among genotypes present in cluster. The maximum inter-cluster distance was

recorded between Cluster II to cluster V (5.76), followed by cluster V and cluster III (5.71), and cluster II and cluster VI (5.57) which indicated the existence of high genetic diversity among genotypes in these clusters, and therefore crosses between the genotypes of these clusters could yield desirable transgressive segregants for desirable quality traits. The minimum inter-cluster was recorded by cluster IV and cluster III (2.96), followed by cluster IV and cluster II (3.03), cluster II and cluster I (3.36), and cluster VI and cluster VII (3.39).

Cluster means of all the quality traits are presented in Table 3. The cluster mean showed different values for all the eleven traits studied. Cluster III had shown the highest mean value for GL (10.05), KL (7.41), LBBC (3.58), KLAC (9.52). The desirable intermediate range of Amycnt (21-25%) was reported by cluster I (23.13), cluster III (23.44) and cluster IV (22.71). The lower value is desirable for grain breadth, kernel breadth, kernel breadth after cooking and elongation ratio

and has been found by cluster III (2.32), cluster III (2.09), cluster IV (2.33) and cluster III (1.27) respectively. "Selection of parents from genetically homogeneous clusters should be avoided to maintain relatively broad genetic base. The maximum amount of desired heterosis will be manifested in cross combinations involving the parents belonging to most divergent clusters for quality characters and is important objective for a plant breeder" [9]. By considering this, genotypes belonging to Cluster II to cluster V, followed by cluster V and cluster III, and cluster II and cluster VI, can be used for making crosses. Cluster III has shown desirable estimates for different quality traits. Naik *et al.*, 2021 studied the experiment with combination of physical (KL, KB, KLAC, KBAC, EI, ER) and physiochemical quality traits (ADV and amyln) and suggested more divergent cluster for breeding program.. Simultaneous consideration of quality traits along with yield contributing traits for cluster analysis were done by Devi *et al.* [10], [11] and Krishna Veni *et al.* [12].

Table 1. Cluster composition of forty rice genotypes

Cluster	Number of Genotypes	Name of Genotypes
1	9	BD105, Swarna, Karhani, URG-22, Sambhamahsuri, HUR3022, HUR105, URG-30, DRR Dhan 48
2	5	Dudhkander, URG-19, IR 85850-AC 157-1, IR 91143-AC 293-1, URG-24
3	13	IR 95133:1-B-16-14-GBS-P1-2-2, IR 95133:1-B-16-14-10-GBS-P5-2-3, IR 95133:1-B-16-14-GBS-P1-2-3, IR15M1633, IR 82475-110-2-2-1-2, IR15M1689, IR64, Sathi, IR 95133:1-16-14-10-GBS-P6-1-5, IR 91143-AC 290-1, IR 91143-AC 239-1, IR15M1546, IR 99642-57-1-1-B
4	7	M-48, M-399, URG-1, BRRIdhan 64, HURZ-3, IR15M1537, BRRIdhan 72
5	2	Nagina-22, BG-102
6	4	IR 95133:1-B-16-14-10-GBS-P5-1-3, HURZ-1, DRR Dhan 45, MTU1010

Table 2. Inter-cluster and Intra-cluster (diagonal) D² of forty rice genotypes

	C1	C2	C3	C4	C5	C6
C1	1.26	3.36	4.11	2.95	4.22	4.40
C2		1.07	3.78	3.03	5.76	5.57
C3			1.06	2.96	5.71	3.59
C4				0.77	3.83	3.39
C5					1.32	3.84
C6						0.92

Table 3. Average performance of clusters for ten quality traits in forty rice genotypes

Cluster	I	II	III	IV	V	VI
GL	8.16	8.54	10.05	8.67	7.63	9.75
GB	2.44	3.09	2.32	2.74	2.84	2.44
KL	5.8	5.81	7.41	6.3	5.59	7.17
KB	2.13	2.28	2.09	2.37	2.43	2.09
LBBC	2.87	2.57	3.58	2.68	2.32	3.45
KLAC	7.87	8.39	9.52	8.41	7.70	9.37
KBAC	2.37	2.75	2.33	2.57	2.85	2.28
EI	1.23	1.20	1.16	1.24	1.18	1.21
ER	1.37	1.43	1.27	1.34	1.38	1.31
ADV	2.23	3.53	4.06	4.18	3.17	3.56
Amycnt	23.13	25.11	23.44	22.71	19.77	20.33

(GL: grain length, GB: grain breadth, KL: Kernel length, KB: Kernel breadth, KLAC: Kernel length after cooking, KBAC: Kernel breadth after cooking. EI: Elongation Index, ER: Elongation ratio, ADV: Alkali digestion value, Amycnt: Amylose content)




Fig. 1. Dendrogram showing relationship among 40 rice (*Oryza sativa* L.) genotypes in six clusters based on Mahalanobis' D^2

3.2 Principle Component Analysis

PCA was used to access the variation and relationships among the eleven studied characters. Almost 90% of variability present among the 40 rice genotypes was captured by the five Principal Components (PC) (Table 4). PC1 contributed for 44.98% of the total variation with LBBC (18.34) having the highest and Amylcnt (0.28) with the lowest loading. PC2 accounted 16.28% of the total variation with KB (22.73) having the highest and ER (0.10) with the lowest loading.

Furthermore, 11.10% variation was explained by PC3 with EI (46.87) and KL (0.02), having

highest and lowest loading, respectively. The remaining two PC, i.e., PC4 and PC5 were captured 10.31% and 7.12% variability. The PC4 has highest loadings for amylnct (25.89) while, the lowest loadings for KBAC (0.29). Similarly, in the case of PC5, amylnct (44.61) had the highest loadings and LBBC has zero percent loading. The variables included in the first PC which explained 44.98% of total variance thus showed high importance for primary selection in under-studied rice breeding lines. Ashok et al. [13], Naik et al. [9] demonstration of the use of factor analysis for efficient selection criteria in rice breeding mprogram provides strong support for our findings.

Table 4. Eigenvalues, percent variance, cumulative variance percent and estimated compound matrix in principle components

Source	PC1	PC2	PC3	PC4	PC5	PC6
Eigenvalue	4.95	1.79	1.22	1.13	0.78	0.57
Variance percent	44.98	16.28	11.10	10.31	7.12	5.20
Cumulative variance percent	44.98	61.26	72.36	82.67	89.79	94.98
GL	13.82	9.79	0.87	2.08	2.49	4.77
GB	9.12	8.00	1.72	4.16	0.19	12.68
KL	16.43	6.07	0.02	4.02	0.05	0.01
KB	5.44	22.38	2.52	22.38	0.23	6.46
LBBC	18.34	1.87	1.08	1.87	0.00	1.38
KLAC	12.67	11.14	3.01	0.58	1.16	16.15
KBAC	7.75	17.85	18.00	0.29	0.68	5.37
ER	10.63	0.10	17.04	1.96	5.49	19.52
EI	3.14	0.16	46.87	20.98	0.09	4.60
Amycnt	0.28	11.39	7.37	25.89	44.61	6.32
ADV	2.38	11.25	1.50	15.78	45.01	22.73

(GL: grain length, GB: grain breadth, KL: Kernel length, KB: Kernel breadth, KLAC: Kernel length after cooking, KBAC: Kernel breadth after cooking, EI: Elongation Index, ER: Elongation ratio, ADV: Alkali digestion value, amylnct: Amylose content)




Fig. 2. Principle component analysis biplot traits

(GL: grain length, GB: grain breadth, KL: Kernel length, KB: Kernel breadth, KLAC: Kernel length after cooking, KLAC-1: Kernel breadth after cooking. EI: Elongation Index, ER: Elongation ratio, ADV: Alkali digestion value, Amylcnt: Amylose content)

The LBBC, GL, KL, KB, GB, KLAC, KBAC and ER had higher vector length indicating the presence of large variability, while remaining traits namely, ADV, Amyl.cnt and EI had smaller vector lengths indicating low variability (Fig. 2). Vector LBBC and ER has formed an angle with any other trait. Traits, EI, GB, KBAC, KB and amylcnt these traits positively correlated with each other. While, vector of LBBC and ER, LBBC and EI, KL and ER were diverged and form a large angle (close to 180°), which indicating they were negative correlated. The vector of ER meets with EI, GB, KBAC, KB and amylcnt, also, LBBC meets vectors of KL, GL and KLAC almost at 90° indicating non-significant or low negative association with these traits. Similar experimental findings were reported by Akinol TF et al. [14], and Naik et al. [9].

4. CONCLUSION

Biometrical technique made it easy to measure the magnitude of variability among the breeding materials.

Selecting contrasting parent lines for hybridization as reflected from D^2 statistics and PCA, would ensure greater chances of obtaining high heterotic hybrids and broad spectrum of variability in segregating progenies. Therefore, the genotypes belonging to Cluster II to cluster V, followed by cluster V and cluster III, and cluster II and cluster VI, can be used for making crosses as they have higher mean value for quality traits and higher inter cluster distance for greater diversity. Hence, by considering the criteria of yield performance, these genotypes can be further utilized in breeding programs for consumer preferred traits. The result of PCA indicate that 90% of diversity is apprehended by first five PCs. Also, EI, GB, KBAC, KB and amylcnt these traits positively correlated with each other. While, the traits, LBBC and ER, LBBC and EI, KL and ER were diverged and form a large angle (close to 180°), which is indicating they were negatively correlated. Consideration of this estimates for formulating the desired results in plant breeding will be beneficial for further varietal improvement program.

ACKNOWLEDGEMENT

The authors thankfully acknowledge IRRI-South Asia Hub, Hyderabad for providing the requisite germplasm and Harvest Plus Project of IRRI

funded by IFPRI-CIAT, USA for their support to get this work accomplished.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Sala M, Ananda kumar CR, Geetha S, Gnanamalar RP, Hemalatha G. Variability studies for iron and zinc content on segregating population of rice. *Electron J Plant Breed.* 2013;4(3):1201-4.
2. Juliano BO, Duff B. Rice grain quality as an emerging priority in national breeding programs. In: Grain R, editor *Marketing and quality issues*. Los Banos, Philippines: International Rice Research Institute; 1991;55-64.
3. Oko AO, Ubi BE, Dambaba N. Rice cooking quality and physico-chemical characteristics: a comparative analysis of selected local and newly introduced rice varieties in Ebonyi State, Nigeria. *Food Public Health.* 2012;2(1):43-9.
4. Rohilla R, Singh VP, Singh US, Singh RK, Khush GS. Crop husbandry and environmental factors affecting aroma and other quality traits. In: Sing RK, Sing US, Khush GS, editors. *Aromatic rices*. Oxford and IBH Publishing Co. Pvt. Ltd. India: New Delli. 2000; 201-16.
5. Mohammadi SA, Prasanna BM. Analysis of genetic diversity in crop plants, salient statistical tools and considerations. *Crop Sci.* 2003;43(4):1235-48.
6. Juliano BO. A simplified assay for milled rice amylose. *Cereal Sci Today.* 16. 1971;340:360:334-8.
7. Rao CR. *Advanced statistical methods in biometrical research*. New York: John Wiley & Sons Inc. 1952:236-72.
8. Singh RK, Chaudhary BD. *Biometrical methods in quantitative genetic analysis*. Ludhiana, New Delhi: Kalyani Publishers; 1977.
9. Naik MVK, Pillai MA, Saravanan S. Genetic diversity analysis for yield associated and quality traits in promising rice varieties of Tamil Nadu. *Int J Bio-Resource Stress Manag.* 2021;12(5): 361-9.
10. Devi RK, Parimala K, Cheralu C. Assessment of genetic diversity and other

- genetic parameters in exotic germplasm of rice. Bioin Folet. 2016;13(1B): 190-6.
11. Asante MD, Adjah KL, Annan-Afful E. Assessment of genetic diversity for grain yield and yield component traits in some genotypes of rice (*Oryza sativa* L.). J Crop Sci Biotechnol. 2019;22(2): 123-30.
 12. Krishna Veni B, Vijaya Lakshmi B, Ramana J. Variability and association studies for yield components and quality parameters in rice genotypes. Rice Res. 2013;6(2):16-23.
 13. Ashok S, Jyothula D, Babu DR. Understanding diversity in rice (*Oryza sativa* L.) germplasm using principle component analysis. Agric Res J. 2017;54(2):170.
 14. Akinola TF, Odiyi AC, Fayeun LS, et al. Genetic variability and genetic diversity of 13 upland rice genotypes for agronomic traits and nutritional qualities. J Agric Sci Bot. 2019;3(1):6-11.

© 2022 Khair et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/92540>